

Pseudo-anévrisme de l'artère radiale : une rare complication de la coronarographie par voie radiale

Radial artery pseudoaneurysm: A rare complication after transradial coronary angiography

Salma Charfeddine*, FatenTriki, Rania Hammami, Leila Abid, Dorra Abid, Samir Kammoun

Department of cardiology, HediChaker university hospital, Sfax, Tunisia

Résumé

La voie radiale est actuellement la voie d'abord de première intention au cours des explorations coronaires. Les complications sont rares. Les pseudo-anévrismes de l'artère radiale est une complication extrêmement rare. Nous rapportons le cas d'une patiente ayant présenté un pseudo-anévrisme de l'artère radiale 7 mois après une coronarographie à visée diagnostique. La présentation clinique et les options thérapeutiques ont été discutées.

Mots-clés

Coronarographie, abord radial, complication vasculaire, pseudo-anévrisme

Summary

The radial approach for diagnostic and therapeutic procedures offers clear advantages in terms of vascular complications. The pseudoaneurysm following radial artery access is an extremely uncommon complication. We report the case of a patient who developed radial artery pseudoaneurysm 7-months after coronary angiography intervention. The clinical presentation and treatment options are discussed.

Keywords

Transradial approach, vascular complication, coronary artery interventions

Correspondance

Salma Charfeddine

Cardiology department, Hedi Chaker University Hospital, Sfax, Tunisia

email: selma_charfeddine@yahoo.fr

INTRODUCTION

The transradial approach has recently been recommended as the first choice to reduce access-site vascular complications during percutaneous coronary diagnostic and interventional procedures [1]. It may be preferred in anticoagulated patients to avoid those vascular complications associated with the femoral approach, which are compounded by systemic anticoagulation [2]. The radial artery pseudoaneurysm is an extremely rare complication associated with transradial access [1, 3, 4]. We report a case of radial artery pseudoaneurysm complicating diagnostic coronary angiography in a patient receiving therapeutic oral anticoagulation.

CASE PRESENTATION

A 64-year-old female with history of hypertension, atrial fibrillation and maintained on chronic warfarin therapy after mechanical mitral valve prosthesis developed a severe tricuspid regurgitation associated with right ventricular failure symptoms. Before redux tricuspid surgery, a diagnostic coronary angiography was performed from the right radial artery with a 5Fr system. The procedure was performed afteracenocoumarol had been stopped for 3 days. The InternationalNormalized Ratio (INR) at the time of angiography was 1.6. The diagnostic angiography revealed no-significant atherosclerosis in coronary arteries. A bolus of 5.000 IU unfractionated heparin bolus was administered though the sheath.A J shape guide wire progressed without any difficulty through the radial artery. The vascular sheath was removed just after the procedure and a compressive dressing was performed. The patient represented 7 months later with painful swelling over the radial puncture site (figure 1).



Figure 1: Image of right radial pseudoaneurysm with area of swelling noted at radial access site.

Colour Doppler ultrasonography demonstrated a pseudoaneurysm with partial thrombosis (Figure 2). A conservative therapy with prolonged local compression failed to close it. The patient required surgical correction. She underwent surgical excision of the pseudoaneurysm and suture of the radial artery. She remained well at follow-up after the procedure, with preserved radial arterial pulse and no residual swelling.




Figure 2: Doppler ultrasonography showing a pseudoaneurysm (horizontal arrow)superior to the radial artery (vertical arrow) access site.

DISCUSSION

The transradial approachfor percutaneous coronary interventionssurely offers advantages in terms of access-site vascular complications. Although uncommon, these complications may not be trivial as some of them may cause serious clinical consequences such as hand ischemia, compartment syndrome and arterial rupture[4]. Previous meta-analysis found up to 1.0% rate of radial access site complications including major haemorrhages and a 65% to 80% rate reduction over the transfemoralaccess[5, 6]. In previous large single centre series, radial artery access clinically relevant vascular complications were around 0.4% [7, 8]. Among the patients who underwent transradial procedures in the radial versus femoral access for coronary angiography and intervention in patients with acute coronary syndromes(RIVAL) trial, the only major non haemorrhagic complication was pseudoaneurysm necessitating closure with a 0.2% incidence [6]. Pseudoaneurysm as a transradial access complication is found in the literature with a rate of 0% to 0.5%. In the largest series up to now, this kind of complication has been reported with an incidence of 0.04% [7].The radial pseudoaneurysm is clearly uncommon when compared to transfemoral access, with less significant morbidity. It

was suggested that ongoing systemic anticoagulation may predispose to radial artery pseudoaneurysm formation [1]. Furthermore, the development of radial artery pseudoaneurysm may be related to inadequate compression post procedure or alternatively to delayed bleeding complicating anticoagulation [1]. As reported previously, our patient presented with a typically painful swelling at the site of the radial access. Our patient's INR at the time of the coronary angiography was 1.6 but she received ongoing therapeutic anticoagulation just after the diagnostic procedure. Thereby, delayed bleeding and inadequate compression, may be the potential contributing factors of pseudoaneurysm constitution in this case. The diagnosis of such complication may be done immediately after the procedure or later. In this present example, the patient presented local symptoms 7 months after the procedure. A surgical intervention was required. In fact,

the late diagnosis of the radial artery pseudoaneurysm is significantly associated with the need for surgical correction [4]. The management of this complication includes compression with a view to thrombosis of the false aneurysm, thrombin injection and surgical closure [9, 10]. Although, the surgical treatment is relatively simple as compared to the femoral artery pseudoaneurysm, a prompt diagnosis may decrease significantly the need for surgery [1, 4].

CONCLUSION

Although infrequent, radial-related vascular complications are possible and may not be trivial. A prompt diagnosis of a radial artery pseudoaneurysm using Doppler ultrasonography is essential to avoid further complications and to decrease the need for surgery.

REFERENCES

- Collins N, Wainstein R, Ward M, Bhagwande R, Dzavik V. Pseudoaneurysm after transradial cardiac catheterization: case series and review of the literature. *Catheter Cardiovasc Interv Off J Soc Card Angiogr Interv*. 1 août 2012;80(2):283-7.
- Popovic B, Freysz L, Chometon F, Lemoine J, Elfarra M, Angioi M, et al. Femoral pseudoaneurysms and current cardiac catheterization: evaluation of risk factors and treatment. *Int J Cardiol*. 14 mai 2010;141(1):75-80.
- Suchoń E, Jąkała J, Dykła D, Depukat R, Krochin M, Matysek J. Radial artery pseudoaneurysm as an extremely rare complication associated with transradial catheterisation. *Kardiol Pol*. 2013;71(5):542.
- Burzotta F, Mariani L, Trani C, Coluccia V, Brancati MF, Porto I, et al. Management and timing of access-site vascular complications occurring after trans-radial percutaneous coronary procedures. *Int J Cardiol*. 1 sept 2013;167(5):1973-8.
- Agostoni P, Biondi-Zoccai GGL, de Benedictis ML, Rigattieri S, Turri M, Anselmi M, et al. Radial versus femoral approach for percutaneous coronary diagnostic and interventional procedures; Systematic overview and meta-analysis of randomized trials. *J Am Coll Cardiol*. 21 juill 2004;44(2):349-56.
- Jolly SS, Yusuf S, Cairns J, Niemelä K, Xavier D, Widimsky P, et al. Radial versus femoral access for coronary angiography and intervention in patients with acute coronary syndromes (RIVAL): a randomised, parallel group, multicentre trial. *Lancet Lond Engl*. 23 avr 2011;377(9775):1409-20.
- Burzotta F, Trani C, Mazzari MA, Tommasino A, Niccoli G, Porto I, et al. Vascular complications and access crossover in 10,676 transradial percutaneous coronary procedures. *Am Heart J*. févr 2012;163(2):230-8.
- Hahalis G, Tsigkas G, Kakkos S, Panagopoulos A, Tsota I, Davlourous P, et al. Vascular Complications Following Transradial and Transulnar Coronary Angiography in 1600 Consecutive Patients. *Angiology*. 29 juin 2015;
- Ganchi PA, Wilhelmi BJ, Fujita K, Lee WP. Ruptured pseudoaneurysm complicating an infected radial artery catheter: case report and review of the literature. *Ann Plast Surg*. juin 2001;46(6):647-50.
- Komorowska-Timek E, Teruya TH, Abou-Zamzam AM, Papa D, Ballard JL. Treatment of radial and ulnar artery pseudoaneurysms using percutaneous thrombin injection. *J Hand Surg*. sept 2004;29(5):936-42.